Muscle Biopsy Procedure

Richard J. Piercy MA VetMB MS PhD Dip.ACVIM MRCVS

Comparative Neuromuscular Diseases Laboratory Royal Veterinary College Royal College Street London NW1 0TU

Email: neuromuscular@rvc.ac.uk

Please contact the laboratory W RE R R N \ R Ø V D P S O/HL PD P Dals Ove can only accept pre-booked samples. Samples must be pre-booked at least KRXUV before submitting, however we cannot guarantee to have spaces available within the short time frame. If this is the case we will offer alternative dates for submission.

Indications / possible diagnoses
Exertional myopathies (RER/EPSM etc)

Materials re quired

Muscle Biopsy Method

- 1. Contact the laboratory WR SUH ERRN \RXU VDPSOHV DW OHDV 2rganise same day or overnight courier service prior to sampling.
- For horses with exertional myopathies the semimembranosis is generally chosen; for equine motor neuron disease, biopsy the sacrocaudalis dorsalis (craniolateral to tail head). For other disorders, choose the site based on the most obviously affected muscle or biopsy several regions.
- 3. Sedate horse and prepare skin for sterile surgery.
- 4. A Bergstrom biopsy needle is suitable, but reliable (and usually better) results are obtained with open biopsy.
- 5. Inject subcutaneously up to 10 ml of local anaesthetic, taking care to avoid direct injection into the muscle layer.

6.